

NWA 856 – 320 grams Enriched Basaltic Shergottite



Figure 1: Photograph of NWA 856 (Djel Ibone) kindly provided by Bruno Fectay and Carine Bidaut illustrating thin fusion crust and interior basaltic texture (scale is 1 cm).

Introduction

Jambon *et al.* (2001, 2002) describe the discovery in March 2001 of another shergottite from Morocco. This sample was originally referred to as “Djel Ibone” and is officially known as NWA 856 (Russell *et al.* 2002). The original piece (~ 5 cm) had a thin black fusion crust that was well preserved – see figure 1 in Jambon *et al.* (2002).

This rock is 186 m.y. old and has been in space about 2 m.y.

Petrography

This meteorite is a fine-grained basalt (figure 1) with gray acicular pyroxene phenocrysts up to 12 mm long (Jambon *et al.* 2002). Augite and pigeonite form as separate crystals. Plagioclase (maskelynite) laths are interstitial as is trace merrillite, apatite, pyrrhotite, chromite, Fe-Ti oxides, silica and baddeleyite.

Shock melt pockets are more abundant than in Shergotty or Zagami and this meteorite is highly fractured at all scales. Chennaoui Aoudjehane *et al.* (2005, 2006) studied the high pressure silica phases.

Mineralogical Mode of NWA856

	Jambon <i>et al.</i> 2002
Pyroxene	68 vol. %
Plagioclase	23
Phosphates	1
Oxides	2
Silica	1
Melt Pockets	2

Terrestrial calcite veins cross-cut this meteorite, but terrestrial weathering appears to be at a minimum because Cs, Ba, Sr and U are not elevated (Jambon *et al.* 2002).

Figure 2 shows a large slab of NWA856 with a thin vein of glass and long needles of pyroxene. Photos can also be seen at <http://www.jpl.nasa.gov/snc/nwa856.html>

Mineral Chemistry

Pyroxenes: As in the Shergotty meteorite, augite and pigeonite are present in NWA856 as separate phases, with no pyroxenes of intermediate composition (figure 3). Pyroxenes are zoned; pigeonite ranges from



Figure 2: Close-up photo of slab of NWA856 showing long needles and glass vein..

En₅₉Fs₂₉Wo₁₂ to En₂₆Fs₅₉Wo₁₅ and augite from En₃₆Fs₃₂Wo₃₂ to En₄₈Fs₃₉Wo₁₃.

Maskelynite: Plagioclase has been shocked to maskelynite Ab₄₈Or₂An₅₀.

Phosphates: Both merrillite and Cl-apatite are present. Channon et al. (2011) reported on the abundances of H, F, Cl and S in apatite.

Stishovite: Relative abundant and large euhedral crystals or thin square needles of stishovite have been observed by Raman spectroscopy (Jambon et al. 2002) in melt pockets of NWA 856.

Baddeleyite: Two minute baddeleyite crystals are reported.

Amphibole: Minute amphibole is reported by Jambon et al. (2002) located in melt inclusions in pyroxene cores.

Whole-rock Composition

Table 1 gives the composition of NWA 856 as reported by Jambon et al. (2001, 2002). The REE pattern of




Figure 3: Pyroxene composition diagram for NWA 856 (data replotted from Jambon et al. 2002).

NWA 856 is similar to those of Shergotty and Zagami (figure 4). This is what we call “enriched”. The Ga/Al ratio (4.1×10^{-4}) indicates that this rock is Martian. Weathering does not appear to have left a significant chemical signature in this desert meteorite (Jambon et al. 2002).

Shirai and Ebihara (2008) have also apparently analyzed NWA856 (and other important shergottites).




Figure 4: Normalized rare earth element diagram for NWA 856 compared with that of Shergotty (data from Jambon *et al.* 2002 and Lodders 2000).

Radiometric Isotopes

Brandon *et al.* (2004) presented preliminary data for the Rb-Sr and Sm-Nd isotope systems. They obtained crystallization ages of 150 ± 32 m.y. and 186 ± 24 m.y. respectively (figure 5 and 6). I'm not sure they obtained precise initial Sr and Nd isotope ratios, and I'm not sure why not, but they seem worried about terrestrial contamination.

Misawa and Yamagouchi (2007) obtained an age of 186 ± 12 m.y. by ion microprobe U/Pb dating baddeleyite grains. Again, they emphasize the age determined is "preliminary".

Cosmogenic Isotopes

Berezhnoy *et al.* (2008) reported on ^{10}Be , ^{26}Al and ^{53}Mn activity and calculate an exposure age of 1.5 m.y. They also quote a 2.6 m.y. age by Mathew *et al.* (2003).

Other Isotopes

Jambon *et al.* (2002) reported oxygen isotopes with $\Delta^{17}\text{O} \sim 0.47 \text{‰}$. Rumble and Irving (2009) obtained $\Delta^{17}\text{O} = 0.23 \text{‰}$.

Processing

Figures 7 and 8 shows the detail of the initial sawing of NWA 856.

References for NWA856




Figure 5: Preliminary Rb-Sr mineral isochron for NWA856 (Brandon *et al.* 2004).




Figure 6: Preliminary Sm-Nd mineral isochron for NWA856 (Brandon *et al.* 2004).



Figure 7: End cut of main mass NWA856 (from internet). That black coating is the fusion crust!

Table 1: Chemical composition of NWA 856.

reference weight	Jambon 2001	Jambon 2002 500 mg.	Jambon 2002	Jambon 2002
SiO ₂				
TiO ₂	0.81	0.81	(a)	
Al ₂ O ₃	6.83	6.83	(a)	
FeO	17.8	17.81	(a)	19.97 (c)
MnO	0.49	0.49	(a)	0.54 (c)
CaO	10.2	10.24	(a)	
MgO	9.51	9.51	(a)	
Na ₂ O	1.28	1.28	(a)	
K ₂ O	0.13	0.13	(a)	
P ₂ O ₅				
sum				
Li ppm		4.06	(b)	
Be		0.355	(b)	
F				
S				
Cl				
Sc		55.7	(b)	54.1 (c)
V		295	(b)	
Cr		3361	(b)	3942 (c)
Co		36.3	(b)	43 (c)
Ni	77	77	(b)	85 (c)
Cu		14	(b)	
Zn		59.1	(b)	66 (c)
Ga		14.66	(b)	
Ge				
As				0.18 (c)
Se				
Br				2.64 (c)
Rb		6.24	(b)	8.2 (c)
Sr		48.7	(b)	56 (c)
Y		18.81	(b)	
Zr		62.8	(b)	69 (c)
Nb		3.37	(b)	
Pd ppb				
Ag ppb				< 50 (c)
Sb ppb				14 (c)
Cs ppm		0.43	(b)	0.41 (c)
Ba		41.3	(b)	46 (c)
La	2.16	2.16	(b)	2.34 (c)
Ce		5.49	(b)	6.1 (c)
Pr		0.786	(b)	
Nd		3.88	(b)	3.9 (c)
Sm	3.88	1.5	(b)	1.68 (c)
Eu	1.5	0.582	(b)	0.62 (c)
Gd	0.58	2.51	(b)	
Tb		0.474	(b)	0.48 (c)
Dy		3.12	(b)	
Ho		0.677	(b)	
Er		1.87	(b)	
Tm				
Yb	1.64	1.64	(b)	1.76 (c)
Lu		0.251	(b)	
Hf		1.55	(b)	2.01 (c)
Ta		0.16	(b)	0.23 (c)
W ppb		430	(b)	520 (c)
Au ppb				4 (c)
Th ppm	0.4	0.398	(b)	0.442 (c)
U ppm		0.096	(b)	0.092 (c)

technique: a) ICP/AES, b) ICP/MS, c) INAA



Figure 8: Close up of slab of Djel Ibone illustrating basaltic texture (whose fingers?).